

BOCHNER FORMULA AND BERNSTEIN TYPE ESTIMATES ON LOCALLY FINITE GRAPHS

LI MA

ABSTRACT. In this paper, we study three typical problems on a locally finite connected graph. The first one is to study the Bochner formula for the Laplacian operator on a locally finite connected graph. We use the Bochner formula to derive the Bernstein type estimate of the heat equation. The second is to derive the McKean type eigenvalue estimate for the principal eigenvalue of the Laplacian operator. The last one to obtain global positive solution to porous-media equation via the use of Aronson-Benilan argument.

Mathematics Subject Classification 2000: 05C50, 53Cxx, 35Jxx, 68R10

Keywords: Bochner formula, heat equation, global solution, porous-media equation

1. INTRODUCTION

In this paper, we study three typical problems on a locally finite connected graph. All these questions are related to heat equations. In fact the eigenvalue problems [1] can be considered as the study of solitary solutions to heat equations. We do believe that the study of heat equation on locally finite connected graphs is an important subject as like it is in the Riemannian geometry. We first study the Bochner formula for the Laplacian operator on a locally finite connected graph. Our Bochner formula is new and should be very useful in the study of eigenvalue estimate of the Laplacian operators on graphs. In fact, by invoking the integration by part trick, one may obtain the Reilly formula on locally finite connected graphs. Once we have the Bochner formula, we use it to study the global behavior of the bounded solution to the heat equation. It is quite nature to ask if we can get a Bernstein type estimate for the solutions to the heat equation on locally finite connected graphs. We can obtain this result. The second problem is to derive the McKean type eigenvalue estimate for the principal eigenvalue of the Laplacian operator. Just in the Riemannian geometry case, we need the concept of Cheeger constant on the locally finite connected graphs. In some sense we introduce the point based Cheeger constant. The last question under our consideration is to obtain global positive solution to porous-media equation via the use of Aronson-Benilan argument. This is a hard question

The research is partially supported by the National Natural Science Foundation of China No. 11271111 and SRFDP 20090002110019.

since we may not have the Sobolev compactness imbedding theorem and it is not easy to the global solution from the exhaustion domain method. We can overcome this difficulty by using the Aronson-Benilan type estimate of the bounded solutions to the Porous-media equation. Our main results are (2), theorems 2 4 and 5 below.

Here is the plan of the paper. In section 2 we study the maximum principle of heat equation and give hint to similar result for nonlinear problems. In section 3, we obtain the Bochner formula and the Bernstein type estimate. We study the McKean type estimate of the principal eigenvalue problem in section 4. In the last section 5, we consider the locally bounded global solution to the Porous-media equation.

2. THE MAXIMUM PRINCIPLE

We start from recalling some definitions and the maximum principle for bounded solution to heat equation. Let $G = (V(G), E(G))$ be an infinite, locally finite, connected graph without loops or multiple edges where $V = V(G)$ is the set of vertices of G and $E = E(G)$ is the set of edges. We still write $x \in G$ when x is a vertex of G . We use the notation $x \sim y$ to indicate the edge connects the vertex x to its neighbor vertex y . We equip V with the symmetric weight $\mu_{xy} \geq 0$ associated to the edge $x \sim y$ such that $\sum_{x \sim y} \mu_{xy} > 0$ for each $x \in V$ and we always assume that our edges are unoriented in the sense that $\mu_{xy} = \mu_{yx}$. We call such a graph the short name the weighted graph. Let $d_x = \sum_{x \sim y} \mu_{xy} > 0$.

We define the space of all square summable functions on G ,

$$l^2(V) = \{f : V \rightarrow R; \sum_{x \in V} d_x f(x)^2 < \infty\}$$

with the inner product

$$(f, g) = \sum_{x \in V} d_x f(x)g(x).$$

Define on $l^2(V)$ the Laplacian operator for the function f ,

$$\Delta f(x) = \frac{1}{d_x} \sum_{x \sim y} \mu_{xy} (f(y) - f(x)).$$

and the norm of the gradient of the function f by

$$|\nabla f|^2(x) = \frac{1}{d_x} \sum_{x \sim y} \mu_{xy} (f(y) - f(x))^2.$$

Note that

$$(\Delta f(x))^2 \leq |\nabla f|^2(x).$$

Fix $x_0 \in G$ and let $r(x) = d(x, x_0)$. Let

$$d_{\pm}(x) = \sum_{\{y \sim x; r(y) = r(x) \pm 1\}} \mu_{xy}.$$

be the number of vertices which are 1 step closer or further to x_0 than x . Define the mean curvature $H(x)$ of the sphere of radius $r(x)$ to x_0 by

$$H(x) = \Delta r(x) = \frac{1}{d_x} \sum_{x \sim y} \mu_{xy}(r(y) - r(x)).$$

It can be verified that

$$H(x) = \frac{d_+(x) - d_-(x)}{d_x}.$$

Then we have the following maximum principle ([2][6] [7] [4]).

Theorem 1. *Assume that there exists some $x_0 \in G$ and a constant $C \geq 0$ such that $H(x) \leq C$ on V . Let $u_0(x)$ be any bounded function on G . Then any bounded solution $u(t, x)$ to the heat equation*

$$u_t = \Delta u$$

with initial data $u(0) = u_0$ satisfies

$$\sup_G |u(t, x)| \leq \sup_G |u_0(x)|$$

for every $t \geq 0$.

The proof of the result above is standard, but for completeness, we give the proof.

Proof. Let $M_1 = \sup\{u(t, x); (t, x) \in (0, T) \times V\}$ and $M_2 = \sup\{u_0(x), x \in V\}$. We consider for positive $C > 0$ the function

$$v(t, x) = u(t, x) - M_2 - \frac{M_1}{R}(d(x, x_0) + Ct)$$

If we denote by $B_R = B_R(x_0)$ the ball with radius R and center x_0 , we may conclude $v(t, x) \leq 0$ for $(t, x) \in \{0\} \times B_R \cup [0, T) \times \partial B_R$, which is the parabolic boundary of $[0, T) \times B_R(x_0)$. On $(0, T) \times B_R$, we have

$$v_t \leq \Delta v.$$

From the maximum principle it follows $v(t, x) \leq 0$ which is equivalent to $u(t, x) \leq M_2 + \frac{M_1}{R}(d(x, x_0) + Ct)$. Letting $R \rightarrow \infty$ we obtain $u(t, x) \leq M_2$ on $[0, T) \times V$. Repeating the arguments for $-u$, yields $|u(t, x)| \leq M_2$ on $[0, T) \times V$. \square

By the result above we can derive the uniqueness of bounded solution to the heat equation on V . In fact the claim follows by considering differences of bounded solutions with same initial condition. Actually we can extend the maximum principle to positive solution to the porous media type equation

$$u_t = \Delta \log u, \quad (0, \infty) \times V$$

with bounded initial data. We shall use this fact below, but we leave the proof of it to interesting readers.

3. BOCHNER FORMULA AND BERNSTEIN ESTIMATE FOR HEAT EQUATION

Following the method of Bakry-Emery, we define

$$\Gamma(f, g) = \frac{1}{2} \{ \Delta(fg)(x) - f(x)\Delta g(x) - g(x)\Delta f(x) \}$$

and

$$\Gamma_2(f, g) = \frac{1}{2} \{ \Delta \Gamma(fg)(x) - \Gamma(f, \Delta g)(x) - \Gamma(g, \Delta f)(x) \}.$$

Then, by direct computation,

$$\begin{aligned} \Delta f^2(x) &= 2f(x)\Delta f(x) + |\nabla f|^2(x), \\ \Gamma(f, g)(x) &= \frac{1}{2d_x} \sum_{y \sim x} \mu_{xy} (f(y) - f(x))(g(y) - g(x)), \\ \Gamma(f, f)(x) &= \frac{1}{2} |\nabla f|^2(x), \\ (1) \quad \Gamma_2(f, f)(x) &= \frac{1}{4} |D^2 f|^2(x) - \frac{1}{2} |\nabla f|^2(x) + \frac{1}{2} (\Delta f)^2(x), \end{aligned}$$

where

$$|D^2 f|^2(x) := \frac{1}{d_x} \sum_{y \sim x} \frac{\mu_{xy}}{d_y} \sum_{z \sim y} \mu_{yz} |f(x) - 2f(y) + f(z)|^2.$$

We now compute the Bochner formula for the function f .

$$-\Delta |\nabla f|^2(x) = -|D^2 f|^2(x) + \frac{2}{d_x} \sum_{y \sim x} \frac{\mu_{xy}}{d_y} \sum_{z \sim y} \mu_{yz} (f(x) - 2f(y) + f(z))(f(x) - f(y)).$$

Set

$$I = \frac{2}{d_x} \sum_{y \sim x} \frac{\mu_{xy}}{d_y} \sum_{z \sim y} \mu_{yz} (f(x) - 2f(y) + f(z))(f(x) - f(y)).$$

Note that

$$\begin{aligned} I &= 2|\nabla f|^2(x) + \frac{2}{d_x} \sum_{y \sim x} \mu_{xy} (f(x) - f(y)) \Delta f(y) \\ &= 2|\nabla f|^2(x) + \frac{2}{d_x} \sum_{y \sim x} \mu_{xy} (f(x) - f(y)) (\Delta f(y) \\ &\quad - \Delta f(x)) + \Delta f(x) \frac{2}{d_x} \sum_{y \sim x} \mu_{xy} (f(x) - f(y)) \\ &= 2|\nabla f|^2(x) + 2|\Delta f|^2(x) - 2(\nabla f, \nabla \Delta f)(x). \end{aligned}$$

Then we have the following Bochner formula:

$$(2) \quad -\Delta |\nabla f|^2(x) = -|D^2 f|^2(x) + 2|\nabla f|^2(x) + 2|\Delta f|^2(x) - 2(\nabla f, \nabla \Delta f)(x).$$

We now use this formula to derive the Bernstein type estimate for the bounded solution $f(t, x)$ to the heat equation

$$f_t = \Delta f$$

with initial data f_0 . Using (2) we get that

$$(3) \quad (\partial_t - \Delta) \left(\frac{1}{2} |\nabla f|^2(t, x) \right) = -\frac{1}{2} |D^2 f|^2(t, x) + |\nabla f|^2(t, x) + |\Delta f|^2(t, x).$$

Assume that on G the curvature condition

$$\Gamma_2(f, f) \geq \frac{1}{m}(\Delta f)^2(x) + \frac{k}{2}|\nabla f|^2(x)$$

for some constants $m > 0$, $k \in \mathbb{R}$.

Using (1) we know that

$$\frac{1}{2}|D^2 f|^2(x) \geq (k+1)|\nabla f|^2(x) + \left(\frac{2}{m} - 1\right)(\Delta f)^2(x).$$

Inserting this back to (3) we get that

$$(\partial_t - \Delta)\left(\frac{1}{2}|\nabla f|^2(t, x)\right) \leq -k|\nabla f|^2(t, x) + \left(2 - \frac{2}{m}\right)|\Delta f|^2(t, x).$$

Using

$$|\Delta f|^2(x) \leq |\nabla f|^2(x)$$

we obtain that

$$(\partial_t - \Delta)\left(\frac{1}{2}|\nabla f|^2(t, x)\right) \leq \left(-k\left(2 - \frac{2}{m}\right)_+\right)|\nabla f|^2(t, x).$$

Recall that

$$(\partial_t - \Delta)f^2(t, x) = -|\nabla f|^2(t, x).$$

Then we can choose $\alpha > 0$ such that

$$(\partial_t - \Delta)(t|\nabla f|^2(t, x)) + \alpha f^2(t, x) \leq 0.$$

Using the maximum principle we then have

$$t|\nabla f|^2(t, x) + \alpha f^2(t, x) \leq \alpha \sup_G f_0^2(x)$$

Since $f(t, x)$ is uniformly bounded in t , we know that there exists $t_k \rightarrow \infty$ such that

$$f(t_k, x) \rightarrow f_\infty(x)$$

for each $x \in G$, and $\lim_{t_k \rightarrow \infty} |\nabla f(t_k, x)|^2 \rightarrow 0$, which implies that $f_\infty(x) = \text{const.}$ In conclusion we have the result below.

Theorem 2. *Assume that there exists some $x_0 \in G$ and a constant $C \geq 0$ such that $H(x) \leq C$ on V . Let $f_0(x)$ be any bounded function on G . Then any bounded solution $f(t, x)$ to the heat equation*

$$f_t = \Delta f$$

with initial data $f(0, x) = f_0(x)$ exists globally and there exists $t_k \rightarrow \infty$ such that

$$f(t_k, x) \rightarrow f_\infty(x)$$

where $f_\infty(x)$ is a constant function.

4. MCKEAN TYPE EIGENVALUE ESTIMATE

We now study the McKean type eigenvalue estimate for the principal eigenvalue λ_1 of the Laplacian operator defined above. Let $\lambda_1(R)$ be the first eigenvalue of the Laplacian operator defined on the ball $B_R(x_0)$ with Dirichlet boundary condition on the boundary $\partial B_R(x_0)$. Then $(\lambda_1(R))$ is a decreasing sequence in R and $\lambda_1 = \lim_{R \rightarrow \infty} \lambda_1(R)$. It is not hard to see that λ_1 does not depend on the based point x_0 . Recall that for any finite subset $\Omega \subset G$, we define Ω^c the complement set of Ω in G . We let $\mu(\Omega) = \int_{\Omega} 1 = \sum_{x \in \Omega} d_x$ and define the measure of the edge boundary $\partial\Omega$ by

$$\mu(\partial\Omega) = |\partial\Omega| = \sum_{\xi=xy \in E(G), x \in \Omega, y \in \Omega^c} \mu_{xy}.$$

It can be directly verify [1] that for any f in the finite set Ω ,

$$(4) \quad \int_{\Omega} \Delta f = \sum_{x \in \Omega} \sum_{y \in \Omega^c} (f(y) - f(x)) \mu_{xy}.$$

We also recall the general integration by part formula. Let $\Omega \subset V$ be a finite subset of V . Define

$$U_1(\Omega) = \{y \in V; \text{dist}(y, \Omega) \leq 1\}.$$

Theorem 3. *For any function f, h , we have*

$$(5) \quad \int_{\Omega} (\Delta f, h) = -\frac{1}{2} \sum_{x, y \in U_1(\Omega)} \mu_{xy} \nabla_{xy} f \nabla_{xy} h$$

where $\nabla_{xy} f = f(y) - f(x)$.

Note that the right hand side of (5) can be written as

$$= -\frac{1}{2} \sum_{x, y \in \Omega} \mu_{xy} \nabla_{xy} f \nabla_{xy} h + \sum_{x \in \Omega, y \in \Omega^c} \mu_{xy} h(x) \nabla_{xy} f.$$

We recall the definition of the Cheeger constant $h(\mu)$ [1] [3] based at x_0 defined by

$$h(\mu) = \inf_{\Omega \subset V; x_0 \in \Omega, \mu(\Omega) \leq \frac{1}{2} \mu(V)} \frac{|\partial\Omega|}{\mu(\Omega)}.$$

Then as in the Riemannian geometry case we have

$$\lambda_1 \geq \frac{h(\mu)^2}{2}.$$

Theorem 4. *Assume that $H(x) \geq a > 0$ for some x_0 and some positive constant a . Then $\lambda_1 \geq a^2/2$.*

Proof. Take $f(x) = r(x) = d(x, x_0)$ in (4) on any finite set Ω containing x_0 . Then Using $H(x) \geq a$ we know that the left side is no less than $\int_{\Omega} H(x) \geq a\mu(\Omega)$ and right side is equals to $\mu(\partial\Omega)$, which implies that

$$\mu(\partial\Omega) = \int_{\Omega} H(x) \geq a\mu(\Omega).$$

Hence $h(\mu) \geq a$, which implies our conclusion. \square

5. GLOBAL SOLUTION TO THE POROUS-MEDIA EQUATION

Given any bounded function $u_0 : V \rightarrow \mathbb{R}_+$. We consider the global existence of the positive solution $u(t, x)$ to the porous-media equation

$$(6) \quad u_t = \Delta \log u, \quad \text{in } (0, \infty) \times V$$

with initial data $u(0) = u_0$. Take any finite subgraph $\Omega \subset V$ and we may first consider (6) in $(0, \infty) \times \Omega$ with initial data and boundary condition u_0 . Let $f = \frac{1}{2} \log u$. Then $u = e^{2f}$ and it satisfies the equivalent problem

$$(7) \quad e^f (e^f)_t = \Delta f, \quad \text{in } (0, \infty) \times V.$$

Actually we can get the local in time solution u_{Ω} to (6) (respectively $f_{\Omega} = \frac{1}{2} \log u_{\Omega}$ to (7)) in $(0, T) \times \Omega$ (for some $T > 0$) by using the discrete Morse flow method [5].

Set $S = \Omega$. For $N > 1$ an integer and any $T > 0$, let

$$h = T/N, \quad t_n = nh, \quad n = 0, 1, 2, \dots, N.$$

Assume that we have constructed $f_j \in L^2(S)$, $0 \leq j \leq n-1$, and f_{n-1} is a minimizer of the functional

$$I_{n-1}(f) = \frac{1}{2h} \int_{\Omega} |e^f - e^{f_{n-2}}|^2 dx + \frac{1}{2} \int_{\Omega} |\nabla f|^2 dx$$

on $H = \{f \in L^2(\Omega) \mid f - f_0 = 0, \text{ on } \partial\Omega\}$. Define

$$I_n(f) = \frac{1}{2h} \int_{\Omega} |e^f - e^{f_{n-1}}|^2 dx + \frac{1}{2} \int_{\Omega} |\nabla f|^2 dx$$

on H . It is clear that the infimum is finite and, by applying the Poincaré inequality to $f - f_0$ [1], that any minimizing sequence is bounded in H . By the direct method of the calculus of variations, one concludes that I_n has a unique minimizer f_n in H which satisfies

$$\frac{1}{h} (e^f - e^{f_{n-1}}) e^f = \Delta f$$

along with the uniform energy bound

$$\frac{1}{2h} \int_{\Omega} |e^{f_n} - e^{f_{n-1}}|^2 dx + \frac{1}{2} \int_{\Omega} |\nabla f_n|^2 dx \leq \frac{1}{2} \int_{\Omega} |\nabla f_{n-1}|^2 dx \leq C.$$

We define $f_N(t) \in H^1$ for $t \in [0, T]$ such that, for $n = 1, \dots, N$,

$$f_N(t) = f_n, \quad t \in [t_{n-1}, t_n].$$

We further define, for $n = 1, \dots, N$,

$$\partial_t e^{f_N}(t) = \frac{1}{h}(e^{f_n} - e^{f_{n-1}}), \quad t \in [t_{n-1}, t_n].$$

Taking a subsequence of $\{f_N\}$ that converges in $L_t^\infty H$, one obtains a limit $f \in L_t^\infty H$ that satisfies

$$e^f \partial_t e^f = \Delta f$$

in distribution sense in the domain $\Omega \times (0, T)$.

To get the globally defined solution, we need the linear upper bound for $u_\Omega = e^{2f}$ and we follow from a well-known argument due to Aronson and Benilan.

Let $\lambda > 1$. Define

$$w_\lambda(t, x) = \lambda u_\Omega(\lambda^{-1}t, x).$$

Then $w_\lambda(t, x)$ satisfies (6) in $(0, T) \times \Omega$ with initial data and boundary condition $\lambda u_0(x)$, which is bigger than $u_0(x)$. By using the comparison principle we know that

$$w_\lambda(t, x) > u_\Omega(t, x) \quad \text{in } (0, T) \times \Omega.$$

Set

$$v_\lambda(t, x) = w_\lambda(t, x) - u_\Omega(t, x).$$

Then

$$\frac{\partial}{\partial \lambda} v_\lambda(t, x) \geq 0, \quad \text{in } (0, T) \times \Omega,$$

or equivalently $u \leq t^{-1}u$ for $u = u_\Omega$, which by integration, implies that $u_\Omega(t, x) \leq C(1+t)$ where $C > 0$ is a constant depending only on u_0 . Hence we can extend the solution $u_\Omega(t, x)$ globally. Take $\Omega = \Omega_j$ where $V = \bigcup \Omega_j$, $\Omega_j \subset \Omega_{j+1}$ are exhaustion subgraphs of V . Then we get a sequence of solutions $\{u_j\}$. By taking subsequence we can get a sub-convergence sequence, still denoted by $\{u_j\}$ and a global (locally bounded) solution $u(t, x)$ of (6) with initial data u_0 such that

$$u(t, x) = \lim_{j \rightarrow \infty} u_j(t, x),$$

locally in $(0, \infty) \times V$. In summary, we then have

Theorem 5. *For any bounded function $u_0 : V \rightarrow \mathbb{R}_+$, there exists a global solution to (6) with initial data u_0 .*

The uniqueness question can be considered by using maximum principle but we leave it open to interesting readers.

REFERENCES

- [1] F. R. K. Chung, *Spectral graph theory*, CBMS Lecture Notes, AMS Publication, 1997.
- [2] Jozef Dodzuik, *Maximum principle for parabolic inequalities and the heat flow on open manifolds*, Indiana Univ. Math. J. 32 (1983), 703C716, <http://dx.doi.org/10.1512/iumj.1983.32.32046>. MR 711862 (85e:58140)

- [3] Jozef Dodzuik, *Difference equations, isoperimetric inequality and transience of certain random walks*, Trans. Amer. Math. Soc. 284 (1984), 787C794, <http://dx.doi.org/10.2307/1999107>. MR 743744 (85m:58185)
- [4] L. Ma, X.Y.Wang, *Kato's inequality and Liouville theorems on locally finite graphs*, Science China: Mathematics, 2013, online.
- [5] Li Ma, I.Witt, *discrete Morse flow for the Ricci flow and Porous media equation*, Arxiv.org, 2012
- [6] ANDREAS Weber, *Analysis of the physical Laplacian and the heat flow on a locally finite graph* (2008), arXiv:0801.0812v2 [math.SP].
- [7] RADOS W. Wojciechowski, *Heat Kernel and Essential Spectrum of Infinite Graphs*, Indiana University Mathematics Journal, Vol. 58, No. 3 (2009)1419-1441

L. MA, DISTINGUISHED PROFESSOR, DEPARTMENT OF MATHEMATICS, HENAN NORMAL UNIVERSITY, XINXIANG, 453007, CHINA
E-mail address: `lma@tsinghua.edu.cn`